

First-Aid Kits

Offshore Medical Kit

Includes COMPLETE BURN and SUTURE KITS. These are the recommended medical supplies for the long distance offshore cruising vessel without a doctor aboard.
 Healthforce Partners: www.shipmd@healthforcepartners.com
 24Hr. PHYSICIAN ADVICE FOR ILLNESS OR INJURY AT SEA.
 Phone 425-806-5770
 Fax 425-806-5771
 Telex(MCI)6838206-MHS-UW
 VHF/SSB Marine Operator Worldwide

Seaside Marine Int'l Drug Co
595 W. 7th Street Suite #207
San Pedro, Ca 90731
Phone: (800) 832-4363 Fax: (800) 548-5527

Date Created/Modified	Kit ID #:	Rx Items Included
2/25/2000	96	<input checked="" type="checkbox"/>

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
1	Seaside Marine Medical Bag	Each	19x16x11 - Red Canvas - Medical Bag		To Store First Aid Supplies	

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
Eye Ear Nose & Throat (Green Rectangle)						
1	Antibiotic Otic Solution (Generic Cortisporin)	10ml			External Ear Infections	Place 2-4 drops in affected ear(s) 3-4 times daily
1	Antipyrine-Benzocaine Otic Solution (Auralgan)	15ml	Analgesic		To reduce pain and inflammation in the ear.	Warm to body temperature and fill the ear canal with 2 to 4 drops. Moisten cotton pledget with the solution and gently insert into the ear canal. Repeat 3 or 4 times daily. Obtain medical advise by radio before using.
1	Cavity (Temporary Filling) 8gm	Each	Temporary Filling Material		Cavities,Inlays, & Post Crowns	Using a suitable filling instrument place direct into cavity and adapt as necessary. It is not essential to dry the cavity prior to placemant. Temporary filling material hardens by absorption of moisture to give an impermeable seal. It is recommended that masticatory pressure is not applied for approximately 15 minutes. For Provisional fixing od post crowns spread a layer on inside of crown and seat in normal way.
18	Cepacol Throat Lozenges	Each	Sore Throat		For temporary relief of occasional minor sore throat pain and dry scratchy throat.	Dissolve 1 lozenge in the mouth every 2 hours as needed or as directed by a physician or dentist.
2	Eye Cup Plastic	Each			For dispensing eye solutions.	1. Examine carefully for any damage. 2. Wash or sterilize thoroughly before each use.
1	Eye Wash Sterile (Squeeze Bottle)	120ml	Eye Irrigating Solution		Use to cleanse irritated eye prior to applying eye pad.	Point the tip of the applicator downward toward the eye and gently squeeze the plastic bottle to irrigate. Use generously.
1	Gentamicin 0.3% Opth Solution (Garamycin)	5ml	Antibiotic		Antibiotic Eye Solution	Apply one or two drops every four hours into the affected eye
2	Guaitussin DM Cough Syrup (Robitussin DM)	120ml	Non-Narcotic Cough Suppressant		For relieving or preventing coughs	Adult: 2 teaspoonfuls every 4 to 6 hours as needed for cough and congestion. May cause drowsiness and sedation. Should not operate machinery.
1	Oxymetozaline Nasal Spray (Afrin)	15ml	Generic Afrin		Decongestant	Adults and children 6 to under 12 years od age (with adult supervision) tilt the head back and apply 2-3 sprays in each nostril not more often than 10 or 12 hours. Do not exceed 2 applications in any 24 hour period. Children under 6 consult a physician
1	Pseudoephedrine 30mg Tablets (Sudafed)	100	Nasal Decongestant		Temporary relief of nasal or sinus congestion. Do not use if you have heart disease, high blood pressure, diabetes, or thyroid disease. May cause difficulty in urination if you have prostatic enlargement.	Adults: two tablets every 4 to 6 hours, no more than 4 doses per 24 hours. Use with caution in people with high blood pressure or heart disease.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
1	Red Cross Toothache Aid	3.75ml	Toothache Medication		For the temporary relief of throbbing, persistent toothache due to a cavity until a dentist can be seen.	Rinse the tooth with water to remove any food particles from the cavity. Use tweezers to moisten pellet, remove excess, and place in cavity. Avoid touching tissues other than tooth cavity. Do not apply more than four times daily or as directed by a dentist.
1	Sulfacetamide 10% Ophth. Solution (Sulamyd)	15ml			For the treatment of conjunctivitis and other superficial ocular infections due to susceptible microorganisms, and as an adjunctive in systemic sulfonamide therapy of trachoma.	Instill two drops into the conjunctival sac or the affected eye every 2 to 3 hours initially. Dosages may tapered by increasing the time interval between doses as the condition responds. The usual duration of treatment is 7 to 10 days. To avoid contamination, do not touch tip of the container to eye, eyelid or any surface.
1	Swim Ear (Alcohol/Glycerin)	30ml	Ear Water-Drying Aid		Helps dry water in ears	Apply 4 to 5 drops in each ear when water remains after swimming, showering, or bathing.
1	Tetracaine 0.5% Ophth Solution (Pontocaine)	15ml	Ophthalmic Anesthetic (Generic Pontocaine)		To produce surface anesthesia of the eye, if necessary, for removal of a foreign body.	One or two drops instilled in eye two or three minutes before beginning procedure.
1	Ventolin HFA Inhaler 200 Doses (Albuterol)	18gm				

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
Wound Care (Large Red Rectangle)						
1	Band Aid Assorted Sheer	60	Sterile		Skin Dressing	Apply as a dressing over small wounds. Do not touch the part that comes in contact with wound.
1	Band Aid Assorted Woven (015050)	20	Dressing		Skin Dressing	Apply as a dressing over small wounds. Do not touch the part that comes in contact with wound.
1	Bloodstopper Bandage (Sterile)	Each	Trauma Dressing			Apply non-stick surface to wound wrap with gauze and cut excess gauze; or use as a sling to secure a splint.
1	Conforming Gauze Bandage 2"x4.1yds (Sterile)	3	Generic Kling		Stretched Rolled Gauze (Sterile)	1. CLEANSE injured area gently and thoroughly with Sterile Rolled Cotton, Cotton Balls, or Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream. 3. COVER injured area completely with Non-Stick Pad to protect wound and absorb fluids. Wrap Rolled Gauze to secure pads if necessary. OR: 4. TAPE on all four sides to firmly secure Non-Stick Pad and seal out dirt and germs.
1	Conforming Gauze Bandage 3"x4.1yds (Sterile)	3	Generic Kling		Stretched Rolled Gauze (Sterile)	1. CLEANSE injured area gently and thoroughly with Sterile Rolled Cotton, Cotton Balls, or Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream. 3. COVER injured area completely with Non-Stick Pad to protect wound and absorb fluids. Wrap Rolled Gauze to secure pads if necessary. OR: 4. TAPE on all four sides to firmly secure Non-Stick Pad and seal out dirt and germs.
1	Conforming Gauze Bandage 4"x4.1yds (Sterile)	3	Generic Kling		Stretched Rolled Gauze (Sterile)	1. CLEANSE injured area gently and thoroughly with Sterile Rolled Cotton, Cotton Balls, or Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream. 3. COVER injured area completely with Non-Stick Pad to protect wound and absorb fluids. Wrap Rolled Gauze to secure pads if necessary. OR: 4. TAPE on all four sides to firmly secure Non-Stick Pad and seal out dirt and germs.
1	Conforming Gauze Bandage 6"x4.1yds (Sterile)	3	Generic Kling		Stretched Rolled Gauze (Sterile)	1. CLEANSE injured area gently and thoroughly with Sterile Rolled Cotton, Cotton Balls, or Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream. 3. COVER injured area completely with Non-Stick Pad to protect wound and absorb fluids. Wrap Rolled Gauze to secure pads if necessary. OR: 4. TAPE on all four sides to firmly secure Non-Stick Pad and seal out dirt and germs.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
6	Cotton Tipped Applicators Sterile 1/Package	Each	6" long		To apply Medication	Do not reuse
2	Elastic Bandage 2"x4.5yds (Ace)	Each	Generic Ace Bandage		An Easy Guide to Treating Minor Strains and Sprains.	1.Unwind 12 to 16 inches at a time. Let bandage relax before wrapping. Start with unstretched bandage to avoid wrapping too tightly. 2. Wrap desired area, overlapping the previous layer by one-half to one-third of its width. 3. Smooth after each turn. 4. Use clips to fasten. 5. If swelling or discomfort occurs after application, loosen the bandage. Remove the bandage before retiring. For best results, unwrap the bandage carefully. Re-roll the bandage, making sure all wrinkles are smoothed out. This is essential to smooth re-bandaging.
2	Elastic Bandage 4"x4.5yds (Ace)	Each	Sprains		An Easy Guide to Treating Minor Strains and Sprains.	1.Unwind 12 to 16 inches at a time. Let bandage relax before wrapping. Start with unstretched bandage to avoid wrapping too tightly. 2. Wrap desired area, overlapping the previous layer by one-half to one-third of its width. 3. Smooth after each turn. 4. Use clips to fasten. 5. If swelling or discomfort occurs after application, loosen the bandage. Remove the bandage before retiring. For best results, unwrap the bandage carefully. Re-roll the bandage, making sure all wrinkles are smoothed out. This is essential to smooth re-bandaging.
2	Eye Pad Large Sterile	Each	2-1/8x2-5/8		Eye protection	Apply as a dressing over inflamed or injured eye
1	Finger Cots Assorted Sizes Sm,Med,& Large	12	Latex		To protect dressing and/or bandage on the fingers	Place on tip of finger and roll to base of hand
1	Gauze Pads 2"x2" 12ply (Sterile)	20	12 Ply Extra Absorbent Gauze Pads		First Aid Gauze Pads are ideal for cleansing and covering larger cuts, scrapes and minor burns because they are highly absorbent. The pads are designed to pull fluids away from the wound site, and comfortably clean dirt and germs from the injured area.	1. CLEANSE injured area gently and thoroughly with Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream or Ointment. 3. COVER injured area completely with Non-Stick Pads to protect wound and absorb fluids. Wrap Rolled Gauze to secure pads if necessary. 4. Tape on all four sides to firmly secure Non-Stick Pads and help seal out dirt and germs.
1	Gauze Pads 4"x4" 12ply (Sterile)	20	12 Ply Extra Absorbent Gauze Pads		First Aid Gauze Pads are ideal for cleansing and covering larger cuts, scrapes and minor burns because they are highly absorbent. The pads are designed to pull fluids away from the wound site, and comfortably clean dirt and germs from the injured area.	1. CLEANSE injured area gently and thoroughly with Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream or Ointment. 3. COVER injured area completely with Non-Stick Pads to protect wound and absorb fluids. Wrap Rolled Gauze to secure pads if necessary. 4. Tape on all four sides to firmly secure Non-Stick Pads and help seal out dirt and germs.
12	Gloves Exam (Non-Sterile) Large	One Pair - Large	Lightly Powdered		Provides Protective Barriers	Use gloves at all times went working with blood or open wounds.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
1	Hibiclens Solution (Chlorhexidine Gluconate)	120ml	Wound Care		Antiseptic/Antimicrobial Skin Cleanser	See bottle for directions.
24	Povidone Iodine Prep Pad (Betadine)	Each	Topical Antiseptic		Apply full strength to skin or mucosal areas by painting, swabbing, or as a wet soak.	Apply full strength to skin or mucosal areas by painting, swabbing, or as a wet soak.
1	Povidone Iodine Scrub (Betadine)	120ml	Topical Antiseptic		For Preoperative Washing by Operating Personnel:	1. Wet hands with water. Pour about 5cc (1 teaspoonful) of BETADINE Surgical Scrub on the palm of the hand and spread over both hands. Without adding more water, rub the Scrub thoroughly over all areas for about five minutes. Use a brush if desired. Clean thoroughly under fingernails. Add a little water and develop copious suds. Rinse thoroughly under running. 2. Complete the wash by scrubbing with another 5cc of BETADINE Surgical Scrub in same way. WARNING: For External Use Only.
1	Surgitube #2 7/8"x5yds	Each	Gauze Bandage		To bandage fingers	1. Slip applicator over finger. Holding gauze at base of finger, withdraw applicator past fingertip. Give applicator one complete twist/tum. 2. Push applicator back over finger. 3. Withdraw and remove applicator. To hold Surgitube in place, use adhesive tape at base of finger.
1	Tape Adhesive 3 in 1 Waterproof	Each	7/8" 5/8" 1/2" strips in 2" roll		To hold bandage in place	Use as needed
2	Tape Adhesive Waterproof 1/2"x5yds (Non Sterile)	Each			To hold bandage in place.	Use as needed.
1	Triangular Bandage	Each	Size: 35"x 50" x 35"		Arm Sling, Tourniquet	Use as Needed
12	Triple Antibiotic Ointment U/D (Neosporin)	1/32oz	Generic Neosporin		Topical Anti-infective	Apply to wound with each dressing change

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
Suturing (Gold Square)						
1	Alcohol Preps	10	Disinfectant		For proper skin cleaning prior to injection.	Remove pad and Rub skin area briskly with pad.
1	Butterfly Closures Medium (Sterile)	10's	In Place Of Sutures		Skin Closures	1. Remove Facing tabs from closure wings. 2. Apply wing sections on either side of wound with nonstick center directly over the wound. 3. Change dressing as directed by your health care professional, or every three days. If healing does not occur in one week or wound shows unexpected changes consult your health care provider. CAUTION: The packaging of this product contains natural rubber latex which may cause allergic reactions.
2	Conforming Gauze Bandage 2"x4.1yds (Sterile)	Each	Generic Kling		Stretched Rolled Gauze (Sterile)	1. CLEANSE injured area gently and thoroughly with Sterile Rolled Cotton, Cotton Balls, or Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream. 3. COVER injured area completely with Non-Stick Pad to protect wound and absorb fluids. Waap Rolled Gauze to secure pads if necessary. OR: 4. TAPE on all four sides to firmly secure Non-Stick Pad and seal out dirt and germs.
2	Conforming Gauze Bandage 3"x4.1yds (Sterile)	Each	Generic Kling		For securing splints, dressings etc.	1. CLEANSE injured area gently and thoroughly with Sterile Rolled Cotton, Cotton Balls, or Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream. 3. COVER injured area completely with Non-Stick Pad to protect wound and absorb fluids. Waap Rolled Gauze to secure pads if necessary. OR: 4. TAPE on all four sides to firmly secure Non-Stick Pad and seal out dirt and germs.
2	Conforming Gauze Bandage 4"x4.1yds (Sterile)	Each	Generic Kling		For Securing Splints, Dressings Etc.	1. CLEANSE injured area gently and thoroughly with Sterile Rolled Cotton, Cotton Balls, or Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream. 3. COVER injured area completely with Non-Stick Pad to protect wound and absorb fluids. Waap Rolled Gauze to secure pads if necessary. OR: 4. TAPE on all four sides to firmly secure Non-Stick Pad and seal out dirt and germs.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
1	Conforming Gauze Bandage 6"x4.1yds (Sterile)	Each	Generic Kling		For securing splints, dressings etc.	1. CLEANSE injured area gently and thoroughly with Sterile Rolled Cotton, Cotton Balls, or Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream. 3. COVER injured area completely with Non-Stick Pad to protect wound and absorb fluids. Waap Rolled Gauze to secure pads if necessary. OR: 4. TAPE on all four sides to firmly secure Non-Stick Pad and seal out dirt and germs.
1	Face Mask W/Shield	Each	For Single Use		Surgical Mask W/faceshield	
1	Forceps Hemostatic Mosquito 5" (Straight)	Each	Stainless Steel			Various uses
1	Forceps Needle Holder Mayo Hegar 5-1/2"	Each	Stainless Steel		To Hold Needle when Suturing.	Use as needed
1	Forceps Splinter 3-1/2" (Stainless Steel)	Each	Stainless Steel		Removing foreign objects	use to remove splinters or foreign objects. Do not dig.
1	Forceps Tissue (1x2) Teeth 5-1/2" Stainless Steel	Each	Stainless Steel		To remove gauze and tissue from wound site.	Sterilize before using.
1	Gauze Pads 2"x2" 12ply (Sterile)	10	12 Ply Extra Absorbent Gauze Pads		First Aid Gauze Pads are ideal for cleansing and covering larger cuts, scrapes and minor burns because they are highly absorbent. The pads are designed to pull fluids away from the wound site, and comfortably clean dirt and germs from the injured area.	1. CLEANSE injured area gently and thoroughly with Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream or Ointment. 3. COVER injured area completely with Non-Stick Pads to protect wound and absorb fluids. Wrap Rolled Gauze to secure pads if necessary. 4. Tape on all four sides to firmly secure Non-Stick Pads and help seal out dirt and germs.
1	Gauze Pads 4"x4" 12Ply (Sterile)	10	12 Ply Extra Absorbent Gauze Pads		First Aid Gauze Pads are ideal for cleansing and covering larger cuts, scrapes and minor burns because they are highly absorbent. The pads are designed to pull fluids away from the wound site, and comfortably clean dirt and germs from the injured area.	1. CLEANSE injured area gently and thoroughly with Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream or Ointment. 3. COVER injured area completely with Non-Stick Pads to protect wound and absorb fluids. Wrap Rolled Gauze to secure pads if necessary. 4. Tape on all four sides to firmly secure Non-Stick Pads and help seal out dirt and germs.
2	Gloves Latex Surgical Size 8 (Sterile)	Each	Latex Surgical Gloves		Maintain Sterility	Instructions For Use. Single Use. Sterile. Do not use if individual package is damaged/open. Powdered with absorbable dusting powder, U.S.P.
1	Lidocaine 1%-50ml W/-Five Syringe/Needle 21gx3cc	Each	Local Anesthetic		As a local anesthetic in minor surgical procedures, as a topical anesthetic on broken skin and mucous membranes.	Dosage varies with the procedure and response of the patient. The smallest dose needed to produce the desired effect should be used. (Consult package insert) Inject slowly and with frequent aspiration to guard against intravascular injection.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
1	Povidone Iodine Solution (Betadine)	120ml	Topical Antiseptic		For use whenever topical germicidal action is required. (Do not use on coral cuts)	Apply full strength to skin or mucosal areas by painting, swabbing, or as a wet soak.
3	Povidone Iodine Swabsticks (3 Per Pack) (Betadine)	Each	Topical Antiseptic		For use whenever topical germicidal action is required. (Do not use on coral cuts)	For external use only. Apply to procedure site as needed. Discard after single use.
1	Povidone Scrub Brush (Betadine)	Each	Surgical Scrub Brush/Sponges with nail cleaner.		For Preoperative Washing by Operating Personnel:	1. For rich lather, saturate sponge with water. SQUEEZE and RELEASE. 2. Begin scrub using detergent solution filled sponge on arms and hands. 3. Complete scrub using brush and sponge. (Nail cleaner as required.) Use brush side-to-side for gentle bristle action, lengthwise for firm bristle action.
1	Scalpel Disposable #11 (Sterile)	Each	Surgical Instrument		Surgery	Obtain medical advice by radio before using !!!
1	Scissors Operating Straight Sharp/Sharp 5-1/2"	Each	Stainless Steel		Surgery	For general use Surgery
1	Sodium Chloride 0.9% Irrigation (Bottle) Suturing	500ml	Plastic Pour Bottle		For cleaning/Irrigating of wounds	Use only if solution is clear and container and seal are intact. Disguard of unused portion.
1	Specimen Container Sterile	120ml	Plastic Jar With Lid		For Urine Sample	
1	Staple Remover (Skin)	Each	Disposable		Remove Skin Staples	(1.) Insert both tips of the lower jaw of the remover under the staple.(2.) Lift slightly so that the staple is perpendicular to the plane of the skin.(3.) Countinue slight lifting while gently squeezing the handles together to reform the staple.(4.) Lift the completely reformed staple straight up from the tissue and discard.
1	Stapler Skin Precise (Disposable)	5	Disposabl;e Skin Staple		Suturing	(1.) Approximate and slightly evert skin edges. Hold stapler upright (90 degrees). Place on skin with gentle downward pressure. Increased pressure results in a more deeply embedded staple.(2.) To form staple, press firmly and completely. Excessive downward pressure on the instrument may make staple removal difficult, and may contribute to a poor cosmetic result.(3.) To release device from staple, relax thumb pressure fully.(4.) Lift stapler away from skin in any direction. Always check staple placement and replace improperly implanted staples. Obtain medical advice by radio before using.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
1	Steri-Strip 1/4"x4" (10/Package) R1546	Each	Steri-Strip Skin Closures Are Made Of Porous, Nonwoven Backing that are coated with a pressure sensitive hypoallergenic adhesive. Steri-strip skin closures are reinforced with filaments for added strength.		Skin Closures	Clean and dry skin at least 3 inches around the wound making sure to remove skin oil or exudate. For best results, use a sponge moistened with either alcohol or saline followed by a firm wipe with a dry sponge. A thin coat of compound benzoin tincture applied to the skin and parallel to the wound will increase strip adhesion
1	Sterile Field 18"x26"	Each	Surgical Drape		Poly-Lined Towel Drape (Sterile)	
1	Sterile Field Fenestrated 18"x26"	Each	Surgical Drape		Poly-Lined Towel Drape; with opening (Sterile)	
1	Sutures Nylon 4-0 18" FS-2 Needle (#662G)	Each	Sterile - Complete - Ready To Go		Suturing	Ordinarily, the advice to the attendant regarding suturing of wounds would be--- DO NOT ATTEMPT IT. However, if many days are to elapse before the patient can be seen by a physician, the attendant should be knowledgeable in the various suture procedures and materials. Suturing is a skill that must be learned through doing, under supervision. There are many complications that can occur if suturing is performed by an inexperienced person. Only under extreme circumstances should suturing be attempted unless the attendant has had specific training in wound closure. It would be more appropriate to use adhesive butterfly strips when possible. See Book For Directions.
1	Sutures Nylon 5-0 18" FS-2 Needle (#661G)	Each	Sterile - Complete - Ready To Go		Suturing	Ordinarily, the advice to the attendant regarding suturing of wounds would be--- DO NOT ATTEMPT IT. However, if many days are to elapse before the patient can be seen by a physician, the attendant should be knowledgeable in the various suture procedures and materials. Suturing is a skill that must be learned through doing, under supervision. There are many complications that can occur if suturing is performed by an inexperienced person. Only under extreme circumstances should suturing be attempted unless the attendant has had specific training in wound closure. It would be more appropriate to use adhesive butterfly strips when possible. See Book For Directions.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
1	Sutures Silk 3-0 18" FS-1 Needle #684G	Each	Sterile - Complete - Ready To Go		Suturing	Ordinarily, the advice to the attendant regarding suturing of wounds would be--- DO NOT ATTEMPT IT. However, if many days are to elapse before the patient can be seen by a physician, the attendant should be knowledgeable in the various suture procedures and materials. Suturing is a skill that must be learned through doing, under supervision. There are many complications that can occur if suturing is performed by an inexperienced person. Only under extreme circumstances should suturing be attempted unless the attendant has had specific training in wound closure. It would be more appropriate to use adhesive butterfly strips when possible. See Book For Directions.
1	Syringe 20cc "Suturing Kit" (Sterile)	.	Latex Free		Irrigation	Obtain medical advice by radio before using!
3	Tape Transpore 1"x10yds (Hypoallergenic)	Each	Surgical Tape		For Taping Various Bandages Etc.	Use as needed
1	Tincture of Benzoin Compound	60ml	Skin Protectant		Useful for treating bed sores, fissures or cracks of the lips. It is also used to protect the skin when excessive taping with adhesive is done.	For external use only. Apply to skin as needed.
1	Triple Antibiotic Ointment U/D (Neosporin) 1/32oz	12	Sterile - Disposable - Single Use "ONLY"		Topical Anti-infective	Apply to wound with each dressing change

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
Burns (Orange Square)						
1	Burn Sheet (Sterile) 60"x96"	Each	Dressing		Sterile Field "Disposable"	Sterility Guaranteed unless package is opened, Damaged or Wet. Avoid freezing and excessive heat. If sheet appears damaged dispose of immediately.
1	Conforming Gauze Bandage 2"x4.1yds (Sterile)	2	Generic Kling		Stretched Rolled Gauze (Sterile)	1. CLEANSE injured area gently and thoroughly with Sterile Rolled Cotton, Cotton Balls, or Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream. 3. COVER injured area completely with Non-Stick Pad to protect wound and absorb fluids. Wrap Rolled Gauze to secure pads if necessary. OR: 4. TAPE on all four sides to firmly secure Non-Stick Pad and seal out dirt and germs.
1	Conforming Gauze Bandage 3"x4.1yds (Sterile)	2	Generic Kling		Stretched Rolled Gauze (Sterile)	1. CLEANSE injured area gently and thoroughly with Sterile Rolled Cotton, Cotton Balls, or Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream. 3. COVER injured area completely with Non-Stick Pad to protect wound and absorb fluids. Wrap Rolled Gauze to secure pads if necessary. OR: 4. TAPE on all four sides to firmly secure Non-Stick Pad and seal out dirt and germs.
1	Conforming Gauze Bandage 4"x4.1yds (Sterile)	2	Generic Kling		Stretched Rolled Gauze (Sterile)	1. CLEANSE injured area gently and thoroughly with Sterile Rolled Cotton, Cotton Balls, or Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream. 3. COVER injured area completely with Non-Stick Pad to protect wound and absorb fluids. Wrap Rolled Gauze to secure pads if necessary. OR: 4. TAPE on all four sides to firmly secure Non-Stick Pad and seal out dirt and germs.
1	Conforming Gauze Bandage 6"x4.1yds (Sterile)	2	Generic Kling		Stretched Rolled Gauze (Sterile)	1. CLEANSE injured area gently and thoroughly with Sterile Rolled Cotton, Cotton Balls, or Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream. 3. COVER injured area completely with Non-Stick Pad to protect wound and absorb fluids. Wrap Rolled Gauze to secure pads if necessary. OR: 4. TAPE on all four sides to firmly secure Non-Stick Pad and seal out dirt and germs.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
1	Gauze Pads 2"x2" 12ply (Sterile)	15	12 Ply Extra Absorbent Gauze Pads		First Aid Gauze Pads are ideal for cleansing and covering larger cuts, scrapes and minor burns because they are highly absorbent. The pads are designed to pull fluids away from the wound site, and comfortably clean dirt and germs from the injured area.	1. CLEANSE injured area gently and thoroughly with Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream or Ointment. 3. COVER injured area completely with Non-Stick Pads to protect wound and absorb fluids. Wrap Rolled Gauze to secure pads if necessary. 4. Tape on all four sides to firmly secure Non-Stick Pads and help seal out dirt and germs.
1	Gauze Pads 4"x4" 12ply (Sterile)	15	12 Ply Extra Absorbent Gauze Pads		First Aid Gauze Pads are ideal for cleansing and covering larger cuts, scrapes and minor burns because they are highly absorbent. The pads are designed to pull fluids away from the wound site, and comfortably clean dirt and germs from the injured area.	1. CLEANSE injured area gently and thoroughly with Gauze Pads, mild soap and water. Pat dry. 2. TREAT and protect wounds against infection with medication or First Aid Cream or Ointment. 3. COVER injured area completely with Non-Stick Pads to protect wound and absorb fluids. Wrap Rolled Gauze to secure pads if necessary. 4. Tape on all four sides to firmly secure Non-Stick Pads and help seal out dirt and germs.
5	Gauze Sponges 4"x4" 12 Ply 2/Pack (Sterile)	Each	Extra Absorbent Pads		Absorbs drainage from wound sites.	Clean and Dressing wound site.
2	Gloves Latex Surgical Size 7-1/2 (Sterile)	Each	Latex Surgical Gloves		Maintain Sterility	Instructions For Use. Single Use. Sterile. Do not use if individual package is damaged/open. Powdered with absorbable dusting powder, U.S.P.
1	Scissors Bandage Stainless Steel (5-1/2")	Each	Lister		To cut next to the skin, with out cutting the skin.	Removal of dressing
1	Silver Sulfadiazine 1% Cream (Silvadene)	50gm	Topical Anti-Microbial		For use on second and third degree burns to aid in prevention and treatment of infection. (Do not use if allergic to sulfa drugs)	The burn wounds are to be cleansed and debrided and Silver Sulfadiazine Cream is applied under sterile conditions. The burn areas should be covered with this cream at all times. The cream should be applied once to twice daily to a thickness of approximately 1/16 inch. When necessary, the cream should be reapplied to any areas from which it has been removed by patient activity. Dressings are not required.
1	Sodium Chloride 0.9% Irrigation (Bottle) Burns	500ml	Plastic Pour Bottle		For cleaning/Irrigating of wounds	Use only if solution is clear and container and seal are intact. Disguard of unused portion.
1	Syringe 20cc "Burn Kit" (Sterile) Luer Lock	Each	Latex Free		Irrigation	Obtain medical advice by radio before using!
2	Tape Micropore Paper 2"x10yds (Hypoallergenic)	Each	Surgical Tape		For Taping Various Bandages Etc.	Use as needed
1	Triple Antibiotic Ointment U/D (Neosporin) 1/32oz	10	Sterile - Disposable - Single Use "ONLY"		Topical Anti-infective	Apply to wound with each dressing change
2	Water-Jel Burn Dressing 2"x6"	Each	Relives Pain, Cools And Comforts, Help Prevent Contamination, Non Adherent		Sterile Gel Soaked Burn Dressing	Tear open package, remove burn dressing, apply dressing immediately to burn area, wrap area without applying pressure, leave dressing in place, then seek medical attention.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
1	Water-Jel Burn Dressing 4"x16"	Each	Sterile Gel Soaked Burn Dressing		For Emergency First Aid use on burns prior to seeking medical care, when necessary. This device is to be used for emergency care only as a temporary measure prior to medical attention.	Tear open package, remove burn dressing, apply dressing immediately to burn area, wrap area without applying pressure, leave dressing in place, then seek medical attention.
1	Water-Jel Burn Dressing 4"x4" (Sterile)	Each	Sterile Gel Soaked Burn Dressing		Relieves Pain-Cools and Comforts	instructions for use: 1. Tear package open. 2. Remove burn dressing. 3. Apply dressing immediately to burn area. 4. Wrap area without applying pressure. 5. Seek medical attention.
1	Water-Jel Burn Dressing-Facial 14"x18" (Sterile)	Each	Sterile Gel Soaked Burn Dressing		For Emergency First Aid Use On Burns Prior To Seeking Medical Care, When Necessary. This Device Is To be used for emergency care only as a temporary measure prior to medical attention.	Tear open package, remove burn dressing, apply dressing immediately to burn area, wrap area without applying pressure, leave dressing in place, then seek medical attention.
3	Xeroform Dressing 4"x4" (Sterile)	Each	Petrolatum Dressing		Wound Dressing For Burns.	The petrolatum gauze bandage is applied in at least two layers over the burned surface and an area extending 2" beyond it. The first dressing should be allowed to remain in place, changing only the outer, dry bandage as needed, for at least 10 days unless signs of infection develop after several days, in which case the dressing should be removed and the burn treated as an infected wound. Watch for blueness or coldness of the skin beyond the dressing and loosen the dressing if they appear.
3	Xeroform Dressing 5"x9" (Sterile)	Each	Petrolatum Dressing		Wound Dressing For Burns.	The petrolatum gauze bandage is applied in at least two layers over the burned surface and an area extending 2" beyond it. The first dressing should be allowed to remain in place, changing only the outer, dry bandage as needed, for at least 10 days unless signs of infection develop after several days, in which case the dressing should be removed and the burn treated as an infected wound. Watch for blueness or coldness of the skin beyond the dressing and loosen the dressing if they appear.
Pain Relievers Non-Narcotic (Small Neon Yellow)						
1	Acetaminophen 325mg Tablets (Tylenol)	100	Generic Tylenol		For Moderate Pain & Fever	Adult Dose: Take 2 tablets every 4 to 6 hours as needed for moderate pain or fever.
1	Ibuprofen 200mg Tablets (Motrin)	100	Generic Motrin Or Advil		For pain,fever, joint inflammation	Adults: take one tablet every 4-6 hours. If pain or fever does not respond to 1 tablet, 2 tablets may be used,do not exceed 6 tablets in 24 hours. Take with food or milk.
3	Ketorolac 30mg/2ml Syringe (Toradol)	2ml				

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
Gastrointestinal & Genitourinary (Gold Rectangle)						
1	Calcium Carbonate 500mg Tablets	150				
1	Cimetidine 200mg Tablets (Tagamet)	30	Generic Tagamet		Stomach Acid Reducer	For relief of symptoms, swallow one tablet with a glass of water. For prevention of symptoms swallow one tablet thirty minutes before ingesting food or beverage that causes heartburn. This product can be used up to twice a day. (Up to two tablets in a 24 hour period)
1	Enema Complete (Disposable)	135ml	Generic Fleet		Saline Laxative	Left side position: lie on left side with knee bent and arms at rest. Knee chest position: kneel, then lower head and chest forward until left side of face is resting on surface. Position arms comfortably. Do not use unless directed by a doctor Adult and children 12 years of age or older. No more than once daily. Do not administer to children under 2 years of age.
1	Hydrocortisone 25mg Suppositories	12	Generic Anusol HC		Hydrocortisone Suppositories Are Indicated For Use In Inflamed Hemorrhoids	Detach 1 Suppository From Strip. Remove The Foil Wrapper. Insert 1 Suppository Into The Rectum Twice Daily Morning And Night For Two Weeks
2	Loperamide 2mg Caplets (Imodium)	12	Anti-Diarrheal		To control the symptoms of diarrhea including travelers diarrhea.	Adult Dose: 2 capsules after first loose stool followed by one capsule after each subsequent loose stool, but no more than 4 capsules per day for no more than 2 days.
1	Meclizine 25mg Chewable Tablets (Bonine)	100			For Seasickness	Take one to two tablets once daily, 1 hour before travel starts. Tablets may be chewed or swallowed whole with water. Thereafter, dose may be repeated every 24 hours for the duration of the travel, or as directed by a physician.
6	Metamucil 3.4gm Packets (Psyllium/Sucrose)	Each			Dietary Fiber Supplement	FIBER LAXATIVE: Adults 12 yrs. & older: One packet in 8oz of liquid, up to 3 times daily at the first sign of irregularity. DIETARY FIBER SUPPLEMENT: Adults 12 yrs & older: One packet in 8oz of liquid, 3 times daily. DIRECTIONS: Pour packet contents into an empty glass. Mix this product with at least 8oz of water or other liquid. Taking this product without enough liquid may cause choking.
20	Phenazopyridine 200mg Tablets (Pyridium)	Each	Generic Pyridium		Burning, frequency and urgency associated with bladder infection.	Dose: 200mg every 8 hours, for 2 days. Note: Turns urine orange. Does not cure infection, antibiotic must be used simultaneously.
1	Prochlorperazine 25mg Suppositories (Compazine)	12				
1	Senna Laxative 8.6mg Tablets (Senokot)	100	Natural Vegetable Laxative		Relieves Constipation	Preferably at bedtime. Adults-Two(2) tablets: Maximum 4 tablets twice daily. Children (6-12)-One(1) tablets: Maximum 2 tablets twice daily, or as directed by a physician.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
1	Transderm Scop 1.5mg (Scopolamine) Transdermal	4	Scopolamine		Motion Sickness Prevention Patches	Dosage and Administration: Apply patch every 3 days as needed. 1. Before applying patch, wash and dry hands thoroughly. 2. Use only one patch at a time. 3. Remove and discard the clear plastic backing from the patch. 4. Apply to the hairless area behind one ear at least 4 hours before traveling. 5. Wash and dry hands thoroughly after application.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
Instruments & Medical Appliances (Lrg Red Square)						
1	Airway Kit Berman Type #0-#6 (7-Bag)	Each	Berman Type		Non Sterile Single Use Only	Obtain radio advice before using!
1	Catheter Tray Urethral With 14fr Catheter	Each	Basic Catheterization Tray		Seek Radio Advice	(1.) Peel back protective cover on outer tray.(2.) Position water-resistant underpad under patient's buttocks.(3.) Don sterile gloves, observing aseptic technique.(4.) Position fenestrated drape so opening is over patient's perineal area.(5.) Open packet of provided swab sticks and thoroughly clean patient's periurethral area, using approved technique.(6.) Using provided lubricating jelly, apply generous amount to first 3" of catheter.(7.) Proceed with catheterization, using approved technique. A urine sample may be collected using the provided specimen container. Obtain medical advice by radio before using.
1	Ear Syringe	60ml	Ear Syringe W/Safety Guard			(1.) Flush syringe thoroughly with warm water before each use. (2.) Use only warm tap water or physician recommended solution at about body temperature. (3.) Child should be seated holding basin under ear to catch returning solution. (4.) Holding ear lobe gently between thumb and forefinger, pull back and upward slightly to maximize exposure of ear canal. Direct ear syringe tip toward the inside of the ear canal, not directly back toward the eardrum. (5.) Gently squeeze bulb to allow the solution to flow into ear canal. When ear canal is fully irrigated, hold bulb loosely so the tip does not interfere with water leaving the ear canal.
1	Finger Splint Padded Assorted Sizes	4	Aluminum With Foam Backing		Immobilizing Fractures/dislocations	The Finger should be straightened by grasping the wrist with one hand and applying traction to the finger tip with the other. The finger should be immobilized with a splint by wrapping in a circular motion starting at the finger tip ending at the base of the hand.
1	IV Butterfly Needle W/Tubing 19gx3/4"-12"	Each	Surflo Winged Infusion Set - For IV use PRO# SV*19BLK		IV	Obtain medical advice by radio before using
1	IV Catheter Angiocath 18gx2" #381147 (Sterile)	Each			IV	Obtain medical advice by radio before using
1	IV Solution Set 15 Drops/ml Macro drip (Sterile)	Each	Sterile Nonpyrogenic Fluid Path		IV	Obtain medical advice by radio before using
1	Pen Light W/Batteries 2-AAA (Heavy Duty)	Each	Reuseable Pen Light		Medical Use	Use as needed
1	Safety Pins Large (Stainless Steel)	12	Various Uses		Attachment	Use As Needed.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
1	Sam Splint 4.5"x36" Orange	Each	Aluminum With Foam Backing		Splint For Arms And Legs Etc.	Use As Directed
1	Scalpel Disposable #15 (Sterile)	Each	Surgical Instrument		Surgery	Obtain medical advice by radio before using !!!
1	Sodium Chloride 0.9% IV Injection (Bag)	1000ml	NACL		To replace chloride and sodium loss.	Before administering, obtain medical advice by radio.
1	Sphygmomanometer Aneroid #104	Each	Blood Pressure Unit		To Measure Blood Pressure	To check for manometer accuracy.(1.) Detach gauge from tubing. Hold gauge in a vertical position.(2.) If the dial needle rests within zero indicator on the faceplate, your unit is guaranteed to be accurate.(3.) Should the needle rest outside the zero indicator, the unit should be returned for calibration.
1	Thermometer 86F to 100F Hypotherm	Each	Low Temp Thermometer		To check for Low Temperature in Hypothermia patient.	See Book For Directions: Obtain medical advice by radio (As Soon As Possible).
1	Thermometer Digital	Each				
6	Tongue Depressor 6" (Sterile)	Each	Deposable/Sterile		To depress the tongue for examining the mouth/throat. Also used for splinting	Use as needed
1	Utility Shears	Each	Medicut Shears		To cut the toughest materials	The 320 features razor-sharp, hardened surgical stainless steel blades with one serrated edge which cuts the toughest materials. Contoured polypropylene handles provide maximum control and comfort. Extra large blade rivet with a rated strength of 88kg assures smooth action and rugged durability. Safety bandage tip. 7-1/4" Long.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
Emergency (Small Blue Square)						
1	Diphenhydramine 25mg Capsules (Benadryl)	24	Generic Benadryl		Antihistamine, for allergies and allergic reactions	Adults take 1-2 capsules (25-50mg) 3 to 4 times daily
1	Diphenhydramine 50mg/ml W/ 21gx3cc Injection	5	Generic Benadryl		Antihistamine, for allergies and allergic reactions.	For use only when the oral form is impractical. Adults- 10 to 50mg IV or deep IM. 100mg if required. Maximum daily dose is 400mg. Children-5mg/kg/24 hours, or 150mg/m2/24 hour, in four divided doses,IV or deep IM Maximum daily dose is 300mg. Obtain medical advice by radio before using.
1	Epinephrine 1mg/ml-1ml W/21gx3cc Syringes	5	Sympathomimetic		Relieve respiratory distress of acute asthma, also to provide rapid relief of severe hypersensitivity reactions to drugs,insect bites, and other allergens.	
1	Nitroglycerin 0.4mg Sublingual Tablets	25	Nitroglycerin		For the prevention and treatment of chest pains associated with angina pectoris	Dissolve 1 tablet under the tongue for chest pain, may repeat every 5 minutes up to 3 doses. Obtain medical advise by radio before using.
1	Pocket Mask PTP Flex Mask	Each	Soft Flex Filter Mask W/Clam Shell Case		CPR	INSTRUCTIONS FOR USE: 1. Fully extend the dome of the mask. 2. Insert one-way valve into the 22mm port of the mask. 3. Position the facemask over the victim's nose and mouth. 4. Open the victim's airway in accordance to recommended guidelines for cardiopulmonary resuscitation. 5. Blow through the mouthpiece to inflate the lungs, using accepted standards for mouth-to-mask resuscitation. 6. If foreign matter appears in the mask such as vomitus or blood, remove the mask and clear the patient's airway. Remove the foreign matter from the mask and mouthpiece by shaking the mask. 7. To provide supplemental oxygen, remove the red cap from the oxygen inlet, securely connect the oxygen tubing to the inlet. 8. Discard the complete unit after use, DO NOT REUSE!!!

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
Skin Preparations (Medium Blue Rectangle)						
1	Blistex Lip Medex	8gm	Lip Balm		Chapped Lips, Wind Burn Etc.	Apply to chapped lips
1	Clotrimazole Vaginal Cream (Gyne Lotrimin)	45gm			Cures Most Vaginal Yeast Infections	Fill The applicator and insert one applicator of cream into the vagina, preferably at bedtime. Repeat this procedure daily for 7 consecutive days. For relief of external vulvar itching, squeeze a small amount of cream onto your finger and gently spread the cream onto the irritated area of the vulva. Use once or twice a day for up to 7 days as needed to relieve external vulva itching. The cream should not be used for vulvar itching due to causes other than yeast infection. If your infection isn't gone in seven days, you may have a condition other than yeast infection. Consult your doctor.
12	Hydrocortisone 1% Cream Unit Dose 1.5gm	Each				
1	Petrolatum Ointment	30gm	Vaseline		For the temporary protection of minor cuts, scrapes, chafed or chapped skin. Protects chafed skin due to diaper rash and helps protect from wetness.	Apply to affected area as often as necessary. Do not use on wounds or burns
1	Pyrrinil Lice Treatment Shampoo	60ml				
1	Sting-Kill Swabs Disposable	5	Contains Benzocaine		For the fast temporary relief of pain and itching due to insect bites, stings and jellyfish.	Apply to the affected area as often as needed but not more than 4 times daily. 1. BEE STINGS ONLY: remove stinger before treatment. 2. Open dispenser cap, then dab moist top with gentle circular rubbing motion on affected area. 3. Replace cap.
1	Tolnaftate 1% Cream (Tinactin)	15gm	Antifungal Cream		Proven clinically effective in the treatment of -athlete's foot - jock itch - ringworm	Wash the affected area and dry thoroughly. Apply a thin layer over the affected area twice daily (morning and night) or as directed by a doctor. Supervise children in the use of this product Do not use in children under 2 years of age. For external use only. Avoid contact with the eyes.
1	Zinc Oxide Ointment 20%	30gm	Skin Protectant		Topical Protective Astringent	Apply as needed in a thin layer to skin areas to be treated or protected.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
RX & Controlled Meds (Medium Dark Green Rectangle)						
1	Acetaminophen W/Codeine 30mg Tablets	100	Analgesic		See packing insert or first aid book for information.	Adult: 1/2 to 1 tablet every 6 hours if needed for pain. Do not take with alcohol. Obtain medical advice by radio before using.
1	Amoxicillin 250mg Capsules	100	Broad-Spectrum Antibiotic		Gram positive bacterial infections. Gram positive (pneumococci, streptococci, and Staphylococci) are organisms which usually cause diseases such as pneumonia, infections of the tonsils and throat, venereal disease, meningitis and septicemia.	Adult dose: 250 to 500mg (one to 2 capsules) every 8 hours for 5 to 10 days. Obtain medical advise by radio before using.
1	Cephalexin 250mg Capsules (Keflex)	100	Antibiotic		Antibiotic	Adult Dose: One to two Capsules 250mg - 500mg (2 to 4 times daily) for 7 to 10 days. Obtain medical advise by radio before using.
1	Diazepam 5mg Tablets (Valium)	100	Sedative		Tranquilizer for tension/nervousness	Adult: Take 1-2 tablets 2-4 times daily, depending upon the condition being treated. Do not take with alcohol. Obtain medical advice by radio before using
3	Diazepam 5mg/ml Carpuject Syringe (Valium)	2ml				
1	Doxycycline 100mg Tablets (Vibramycin)	50	Antibiotic		Upper and lower respiratory bacterial infections, Urethritis, Prostatitis, Epididymitis.	Dose: 100mg every 12 hours for 10 days. Note: Avoid prolonged exposure to direct sun light or crew member may wear sun block. May cause heart burn, do not take before sleeping. Avoid taking if allergic to tetracycline. Take 1-2 hours before eating
1	Erythromycin 250mg Tablets	100	Bacteriostatic Antibiotic (Alternate For Penicillin)		For Infections Of The Upper And lower Respiratory Tract; Infections of the mouth, gums, and teeth; infections of the nose, ears, and sinuses. Can also be effective against amoebas of the intestinal tract which cause amoebic dysentery.	Adult dose: One to 2 tablets (250 to 500mg) every six hours for 7 to 10 days
1	Prednisone 5mg Tablets	100	Prednisone tablets contain prednisone which is a glucocorticoid. Glucocorticoids are adrenocortical steroids, both naturally occurring and synethetic that are readily absorbed from the gastrointestinal tract		Endocrine disorders,rheumatic disorders,collagen diseases,dermatologic diseases,allergic states,ophthalmic diseases, respiratory diseases, hematologic disorders,neoplastic diseases, gastrointestinal diseases.	Dosage of Prednisone tablets should be individualized according to the severity of the disease and the response of the patient. Notify doctor by radio before using.

Qty	Product Name	Size	Description	Off Shore	General Uses	Directions
1	Sulfamethoxazole/Trimeth 400mg/80mg Tablets	100	Urinary Anti- infective (Generic Bactrim S/S)		Acute. Recurrent, or chronic urinary tract infections, primarily cystitis, pyelitis, and pyelonephritis due to organisms susceptible to the drug. Use for treatment of unresponsive traveler's diarrhea.	Adults: 2 tablets twice a day. Take with plenty of fluids. May sensitize skin to sunlight. Do not take if allergic to sulfa drugs. Sore throat, fever, unusual bleeding or bruising, rash, and feeling tired are early signs of serious blood disorders. Obtain medical advice by radio before using !!!
50	Theophylline ER 200mg Tablets (Theolair)	Each	Generic Theolair		Wheezing associated with asthma, lung infections and emphysema. Use with caution in crew members with heart and liver disease. May cause nausea, vomiting and rapid heart rate.	Take 200mg orally every 8-12 hours.